



Unit 1: Number Sets & Exponents**Pythagorean theorem**, to find:

Hypotenuse: $c = \sqrt{a^2 + b^2}$
 Ex: $x = \sqrt{5^2 + 12^2}$
 $x = \sqrt{169} = 13$



Leg: $a = \sqrt{c^2 - b^2}$
 Ex: $x = \sqrt{29^2 - 21^2}$
 $x = \sqrt{400} = 20$


**Changing repeating decimal to fraction:**

Ex: $0.\overline{16} \rightarrow 1.\overline{6} = 1\frac{6}{9} = 1\frac{2}{3} = \frac{5}{3} \rightarrow \frac{5}{30} = \frac{1}{6}$

$\in$: element of

$\subseteq$: subset of

$N \subseteq Z \subseteq Q \subseteq R$

**Laws of exponents:**

$A^0 = 1$ $a^1 = a$

Law	Example
1. $a^m a^n = a^{m+n}$	$x^2 x^3 = x^5$
2. $\frac{a^m}{a^n} = a^{m-n}$ if $m > n$	$\frac{x^6}{x^2} = x^4$
$= \frac{1}{a^{n-m}}$ if $n > m$	$\frac{x^2}{x^6} = \frac{1}{x^4}$
3. $(a^m)^n = a^{mn}$	$(x^2)^3 = x^6$
4. $(ab)^n = a^n b^n$	$(2x)^3 = 8x^3$
5. $\left(\frac{a}{b}\right)^n = \frac{a^n}{b^n}$	$\left(\frac{2}{x}\right)^3 = \frac{8}{x^3}$
6. $a^{-n} = \frac{1}{a^n}$	$x^{-3} = \frac{1}{x^3}$
7. $a^{1/n} = \sqrt[n]{a}$	$x^{1/3} = \sqrt[3]{x}$

Scientific notation

$a \times 10^n$
 $(1 \leq a < 10)$

Ex: $(2.3 \times 10^{-3})(7.9 \times 10^5)$
 $= (2.3)(7.9) \times (10^{-3+5})$
 $= 18.17 \times 10^2$
 $= 1.817 \times 10 \times 10^2$
 $= 1.817 \times 10^3$

Unit 2: Algebraic Expressions**Definitions:**

1. Monomial: 1 term either variable, constant, or product
2. Polynomial: more than one term separated by $+$ or $-$; all terms have variables with natural exponents
 - (a) Binomial- 2 terms
 - (b) trinomial- 3 terms
3. Coefficient: is the number preceding the variable(s)
4. Like terms: have identical variables with identical exponents
5. Degree: (a) of a monomial: sum of exponents of all variables
 (b) of a polynomial: the degree of the term with highest degree
6. Zero of a polynomial: is the value of the variable that makes the polynomial = 0
7. Simplify: is to collect (add or subtract) all like terms to have fewer terms
8. Evaluate: is to replace the variable with a given value & follow BEDMAS

Adding/Subtracting Polynomials: coefficients only, exponents don't change

Ex 1: $(2x^2 + 5x + 8) + (x^2 - 4x + 5)$
 $= 3x^2 + x + 13$

Ex 2: $(2x^2 + 5x + 8) - (x^2 - 4x + 5)$
 $= x^2 + 9x + 3$

Multiplying Polynomials:

Case 1 Distributive	Case 2 FOIL	
Ex: $3x(4x + 2)$ $= 12x^2 + 6x$	Ex: $(3x + 2)(4x - 5)$ F O I L $= 12x^2 - 15x + 8x - 10$ $= 12x^2 - 7x - 10$	Ex: $(3x + 2)^2$ $= (3x + 2)(3x + 2)$ $= 9x^2 + 6x + 6x + 4$ $= 9x^2 + 12x + 4$

Dividing Polynomials: Divide each term by the monomial

Ex: $\frac{20xy^5 - 15xy^2 + 30x^2y^4 + 5xy}{5xy}$
 $= 4y^4 - 3y + 6xy^3 + 1$

Greatest Common Factor:

1. Find the GCF for the coefficients and each variable (the variables with the smallest exponents)
2. To get the second factor, divide the polynomial by your GCF

Ex: $18x^4y + 12x^3y^2$
 $= (6x^3y)(\frac{18x^4y + 12x^3y^2}{6x^3y})$
 $= (6x^3y)(3x + 2y)$

Unit 3: Equations & Inequalities

	Equations	Inequalities/ interval solution	EFF
1	$x + 3 = 7$ $x = 4$	$x + 3 < 7$ $x < 4$	Interval: $]-\infty, 4[$
2	$2x + 4 = 10$ $2x = 6$ $x = 3$	$2x + 4 \geq 10$ $2x \geq 6$ $x \geq 3$	Interval: $[3, \infty[$
3	$5x + 25 = -3x - 23$ $8x + 25 = -23$ $8x = -48$ $x = -6$	$5x + 25 > -3x - 23$ $8x + 25 > -23$ $8x > -48$ $x > -6$	Interval: $]-6, \infty[$
4	$6(x - 2) = -4(2x + 1)$ $6x - 12 = -8x - 4$ $14x - 12 = -4$ $14x = 8$ $x = \frac{8}{14} = \frac{4}{7}$	$6(x - 2) \leq -4(2x + 1)$ $6x - 12 \leq -8x - 4$ $14x - 12 \leq -4$ $14x \leq 8$ $x \leq \frac{8}{14} = \frac{4}{7}$	Interval: $]-\infty, \frac{4}{7}]$
5	$\frac{3x+5}{4} = \frac{4x+10}{2} + 1$ $\frac{3x+5}{4} = \frac{2(4x+10)}{4} + \frac{4}{4}$ $\frac{3x+5}{4} = \frac{8x+20}{4} + \frac{4}{4}$ $3x + 5 = 8x + 20 + 4$ $-5x + 5 = 24$ $-5x = 19$ $x = -\frac{19}{5}$	$\frac{3x+5}{4} \leq \frac{4x+10}{2} + 1$ $\frac{3x+5}{4} \leq \frac{2(4x+10)}{4} + \frac{4}{4}$ $\frac{3x+5}{4} \leq \frac{8x+20}{4} + \frac{4}{4}$ $3x + 5 \leq 8x + 20 + 4$ $-5x + 5 \leq 24$ $-5x \leq 19$ $x \geq -\frac{19}{5}$	Interval: $[-\frac{19}{5}, \infty[$ Exception: when you divide by a negative, flip the sign.

More examples:

$-5(2x+1) + 3(x-2) = 2(4x-1) - 3(2x-3)$
 $-10x - 5 + 3x - 6 = 8x - 2 - 6x + 9$
 $-7x - 11 = 2x + 7$
 $-9x - 11 = 7$
 $-9x = 18$
 $x = -2$


$5(2x+1) + 3(x-2) \geq 2(4x-1) - 3(2x-3)$
 $-10x - 5 + 3x - 6 \geq 8x - 2 - 6x + 9$
 $-7x - 11 \geq 2x + 7$
 $-9x - 11 \geq 7$
 $-9x \geq 18$
 $x \leq -2$

$80 < 4x + 20 < 100$

$60 < 4x < 80$

$15 < x < 20$

Interval solutions is: $]15, 20[$

**Unit 4: Functions**

A relation:		It is a function if there is at most one y-value for every x-value.
Input	Output	
Source	Target	
Antecedent	Image	
Independent	Dependent	
x-value	y-value	
Domain	Range	

Notation: $f(x) = y$

Modes of representation:

1. Written description EX: Repairman charges \$30 per hour plus \$60 for travel.
2. Rule/Equation: $f(x) = 30x + 60$; indep. var. (x) time; dep var. (y) cost
3. Table of values:

X(hours)	0	1	2	3
Y(cost)	60	90	120	150
4. Graph:

Rate of change:

ROC = $\frac{\text{rise}}{\text{run}}$ (given graph)

$= \frac{\Delta y}{\Delta x}$ (given TOV) EX: $\frac{\Delta y}{\Delta x} = \frac{30}{1} = 30$

$= \frac{y_2 - y_1}{x_2 - x_1}$ (given 2 points) EX: A(3,6) and B(5,-2) ROC = $\frac{-2-6}{5-3} = \frac{-8}{2} = -4$

Finding the rule/equation of the line: $y = ax + b$

EX:

X	Y
5	280
8	406

Step 1: find a (ROC)

$a = \frac{406-280}{8-5} = \frac{126}{3} = 42 \text{ \$/hr}$

Step 2: find b (initial value)

$b = y_1 - ax_1$
 $= 280 - (42)(5) = 70\$$

Therefore: $f(x) = y = 42x + 70$

Types of functions:


Constant :	Direct linear:	Partial linear:	Rational:
$y = b$ 	$y = ax$ a: ROC b = 0 	$y = ax + b$ a: ROC b: y-intercept or initial value 	$y = \frac{c}{x}$ $x \cdot y = C$ Constant product

Solving a system of 2 equations:

$\begin{cases} y_1 = 2x + 5 \\ y_2 = x + 8 \end{cases}$ $\begin{cases} y_1 = y_2 \\ 2x + 5 = x + 8 \\ x = 3 \end{cases}$ $\begin{cases} y_1 = 2(3) + 5 \\ = 11 \end{cases}$ $\begin{cases} y_2 = (3) + 8 \\ = 11 \end{cases}$

Therefore the solution is (3,11) : for $x = 3$ the $y = 11$ for both equations

$P = 4w$ $A = w^2$	$P = 2L + 2W$ $A = LW$	$P = 2L + 2W$ $A = LH$	$P = b + s_2 + s_3$ $A = \frac{bh}{2}$
$C = \pi d = 2\pi r$ $A = \pi r^2$	$P = B + b + s_3 + s_4$ $A = \frac{(B+b)h}{2}$	$P = 4w$ $A = \frac{Dd}{2}$	$P = nw$ $A = \frac{pa}{2}$

Unit 5: Geometry**Views:**

A prism: has only 2 parallel and congruent bases, its named after the base, its height is the distance between the bases.

A cylinder: has only 2 parallel and congruent disks, its height is the distance between the bases.

A Pyramid: has 1 base and an apex, its named after the base, it has a height and a slant height $s^2 = h^2 + a^2$ or $s = \sqrt{h^2 + a^2}$ a is apothem

A cone: has one circle base and an apex, has a height and a slant height, $s^2 = h^2 + r^2$


A sphere: a ball, no edges, no vertices, no bases.

Areas And Volumes:

1D: $\times/\div 10$ each time

Area: $\times/\div 100$ each time

Volume: $\times/\div 1000$ each time




Shape	A_L (unit ²)	A_T (unit ²)	VOLUME (unit ³)
Cube	$4w^2$	$6w^2$	w^3
Prism	$P_b h$	$P_b h + 2A_b$	$A_b h$
Cylinder	$2\pi r h$	$2\pi r h + 2\pi r^2$	$\pi r^2 h$
Pyramid	$\frac{P_b s}{2}$	$\frac{P_b s}{2} + A_b$	$\frac{A_b h}{3}$
Cone	$\pi r s$	$\pi r s + \pi r^2$	$\frac{\pi r^2 h}{3}$
Sphere	$A_{LAT} = A_{TOT} = 4\pi r^2$		$\frac{4\pi r^3}{3}$
Hemisphere	$2\pi r^2$	$3\pi r^2$ If base is included	$\frac{2\pi r^3}{3}$

P_b : Perimeter of a base; A_b : Area of base; r : radius; s : slant height; h : height

EX: Find the total surface area and volume of this solid

$$\begin{aligned}
 A_T &= A_{Lcone} + A_{Lcylinder} + A_{Lhemisphere} \\
 &= \pi r s + 2\pi r h + 2\pi r^2 \\
 &= \pi(3)(5) + 2\pi(3)(8) + 2\pi(3)^2 \\
 &= 15\pi + 48\pi + 18\pi \\
 &= 81\pi \text{ cm}^2 \text{ or } 254.47 \text{ cm}^2
 \end{aligned}$$

$$\begin{aligned}
 V &= V_{cone} + V_{cylinder} + V_{hemisphere} \\
 &= \frac{\pi r^2 h}{3} + \pi r^2 h + \frac{2\pi r^3}{3} \\
 &= \frac{\pi(3)^2(4)}{3} + \pi(3)^2(8) + \frac{2\pi(3)^3}{3} \\
 &= 12\pi + 72\pi + 18\pi \\
 &= 102\pi \text{ cm}^3 \text{ or } 320.44 \text{ cm}^3
 \end{aligned}$$

**Unit 6: Missing Measures**

1. EX: A square base pyramid has volume 223 cm^3 , and height 12 cm, find the side length of its base.

$$\begin{aligned}
 V_{pyr} &= \frac{A_b h}{3} \\
 223 &= \frac{A_b 12}{3} \\
 669 &= (s^2)(12) \\
 55.75 &= s^2 \\
 s &= 7.47 \text{ cm}
 \end{aligned}$$

2. Similar solids have a ratio k between them (each dimension of one is k times bigger than the other) it is easier to k

1. Start a table with k, k^2, k^3

2. Solve for k first

3. Set up equal ratios

4. Cross multiply to solve for missing dimensions

EX: Find the

$k = \frac{s'}{s}$	$k^2 = \frac{A'}{A}$	$k^3 = \frac{V'}{V}$
$\sqrt[3]{\frac{125}{8}} = \frac{5}{2}$	$\left(\frac{5}{2}\right)^2 = \frac{25}{4}$	$\frac{125\pi}{8\pi} = \frac{125}{8}$ given

$$k = \frac{r'}{r}$$


$$\frac{5}{2} = \frac{r'}{2}$$

$$r' = 5$$

$$A_b = \pi r^2$$

$$= \pi(5)^2$$

$$= 25\pi \text{ cm}^2$$

**Unit 7: Probability**

Basic counting principle: if there are m ways to do one thing, and n ways to do another, there are $(m)(n)$ ways to do both.

$n!$ ways to arrange all n items


Probability of an event:

$$P(A) = \frac{\text{\# of desired outcomes}}{\text{total \# of outcomes}}$$

Geometric probability:

- 1D: $P(\text{target}) = \frac{\text{target length}}{\text{total length}}$
- 2D: $P(\text{target}) = \frac{\text{target area}}{\text{total area}}$
- 3D: $P(\text{target}) = \frac{\text{target volume}}{\text{total volume}}$

EX: What is the probability of hitting the black?



$$\begin{aligned}
 P(B) &= \frac{A_{\text{black circle}} - A_{\text{white circle}}}{A_{\text{square}}} \\
 &= \frac{\pi R^2 - \pi r^2}{s^2} \\
 &= \frac{\pi(3)^2 - \pi(2)^2}{6^2} \\
 &= \frac{9\pi - 4\pi}{36} \\
 &= \frac{5\pi}{36} = 0.436
 \end{aligned}$$

Unit 8: Statistics**Type of survey:**

CENSUS: whole population; **POLL:** small sample; **STUDY:** Experts in topic

Type of variable/data: 1-**QUALITATIVE** (quality/ non-numerical); or 2-**QUANTITATIVE**(numerical): **Discrete** (Integers); or **Continuous** (Real #s in intervals)

Mode(Mo): data value with the highest frequency

Median(Md): data value at the center of an ordered list

Mean($\bar{x}$): average of all values of data

EX:

$$Mo = [100, 110];$$

Md = the tenth class:

$$[110, 120];$$

$$\bar{x} = \frac{2215}{19} = 116.58$$

Height	Freq.	Midpoint	Total height
[100,110[	8	105	8(105)=840
[110,120[	2	115	2(115)=230
[120,130[	7	125	7(125)=875
[130,140[	2	135	2(135)=270
Total	19		2215

Quartiles: Q1, Q2, Q3 divide the ordered list of data into 4 groups having the same number of data in each. The **Q2** is the median, we find it first.

Range: $R = X_{\max} - X_{\min}$

Interquartile range: $I = Q3 - Q1$

Tables and diagrams:

EX:

1. Bar graph

2. Pie chart

3. Histograms

4. Box and

Whiskers plot.

